



Design and characterization of *swietenia mahagoni jacq.* Mediated silver nanoparticles for hyperglycaemic and hyperlipidaemic control

L.V. Vigneshwaran*

Department of Pharmaceutics, Aditya Bangalore Institute of Pharmacy Education & Research, Yelahanka, Bangalore-560064, Karnataka, India.

ABSTRACT



The present study was designed to evaluate the design and characterization of *swietenia mahagoni jacq.* mediated silver nanoparticles for hyperglycemic and hyperlipidemic control. Using the extraction methods like soxhlet extraction, benzene extraction, Ethanolic Extraction, Ethyl acetate extraction, petroleum ether extraction. Animals were randomly divided into 5 groups. Group 1- normal control, Group 2- diabetic control, Group 3 –treatment control *Swietenia mahagoni* 100mg/kg, Group 4 – *Swietenia mahagoni* 200mg/kg treated group, Group 5 – Glibenclamide 5mg/kg. The levels of biochemical parameter, body weight, serum lipid profile, serum total cholesterol, serum glycerides, VLDL AND HDL cholesterol, HDL cholesterol, Hemoglobin & glycated hemoglobin were examined in each of these groups. Chronic administration of *swietenia mahagoni* at a dose of 100mg/kg & 200mg/kg, p.o. for a period of 21 days markedly decreased the level of blood glucose level & lipid level. Pharmacological studies were assessed for bioassessment, clinical trials, and toxicological studies. Quantitative estimation of total phenolic and flavonoid like estimation of total phenolic content, total flavonoid content, HPTLC investigation of ethanol extract. Taken together, the results suggested that treatment reduced hyperglycemic and hyperlipidemic control. This study demonstrates the *swietenia mahagoni jacq.* Mediated silver nanoparticles for hyperglycemic and hyperlipidemic control.

Keywords: Loratadine; Phenylephrine; Bilayer tablets.

ISSN: 2581-9143

Research Article

Corresponding Author

Name: L.V. Vigneshwaran

Email: vigneshwaran85@gmail.com

Article Info

Received on: 18-10-2025

Revised on: 21-11-2025

Accepted on: 07-11-2025

DOI: <https://doi.org/10.33974/ijrpst.v5i1.239>



Copyright© 2025, L.V. Vigneshwaran., Design and characterization of *swietenia mahagoni jacq.* Mediated silver nano-particles for hyperglycaemic and hyperlipidaemic control, Production and hosting by Rubatosis Publications.

INTRODUCTION

One of the fields that has advanced the fastest in recent years is nanotechnology. The understanding of chemistry, physics, engineering, and material science are all connected in this multidisciplinary field. [1,2] The Greek word "nanos" is where the term "nano" originates, which means "dwarf". [3] In mathematics, a nano is one billionth of a number or period, which is represented as 1×10^{-9} or basically

as 10^{-9} . [4] It is the study of extremely minute things, including almost any science that deals with altering materials at the atomic and molecular scales and comprehending the universe at the atomic level. [5] Nanoparticles, which range in extent from 10 to 1000 nm (1.0 μm), are small solid colloidal particles that contain the active component or organically active substance melted, tricked, adsorbed, or connected. [6] Using a controlled and targeted drug delivery technique, nanotechnology aims to identify as early and accurately as possible and treat effectively without causing any adverse effects. [7] Drug delivery and related pharmaceutical development should be understood as the science in the perspective of nanomedicine, and the technology of multifaceted systems at the nanoscale, having at least two components, one of which is a pharmaceutically active element. [8]

Characterization of Nanoparticles

- Particle size and size distribution: PCS, TEM, SEM, AFM [9]
- Charge determination: Laser Doppler Anemometry, Zeta potentiometer [10]
- Surface Hydrophobicity: Water contact angle measurements, Rose Bengal binding, X-ray photoelectron spectroscopy [11]
- Carrier drug interaction: DSC, FTIR, XRD [12]

- Drug Stability: Bioassay of drug extracted from nanoparticles, Chemical analysis of drug [13]
- Nanoparticle dispersion stability: Critical flocculation temperature [14]
- In vitro drug release: Franz diffusion cell, Dialysis bag method, Open-ended tube method [15]

Preparation of Nanoparticles [16-18]

The three processes for creating nanoparticles are

- Dispersing premade polymers
- Polymerizing monomers
- Co-acervation of hydrophilic polymers or ionic gelation.

The literature has documented various methods for creating nanoparticles making particles with supercritical fluid methods and non-stick molds. It was claimed that the latter had total control over the particles' size, shape, and makeup, and that this may be used as a model for future industrial mass production of NPs.

Drug profile: It's Known as "*Mahogany*," *Swietenia mahagoni* Jacq. The Meliaceae family includes the economically significant, deciduous timber tree, Puerto Rican mahogany, also known as Spanish mahogany, Cuban mahogany, or Jamaican mahogany, native to Central America. One of the most widely used traditional remedies in Africa comes from this important plant, which is closely linked to the African genus *Khaya*.^[19]



Figure 1: Habitat of *S. mahogini jacq*

History: At first, only the British-controlled islands in the West Indies were referred to by the term mahogany. The origin of the name is uncertain, though it may be a distorted form of "Moganwo." The Yoruba and Igbo people of West Africa use this name for trees in the *Khaya* group, which are closely related to *Swietenia* trees. According to the aforementioned interpretations³, the tree's exact origin is still up for debate. The term "mahogany" was first used in print in John Ogilby's *America* in 1671. Although the quality and character of mahogany varied depending on the soil and environment, it was considered a single species until the 19th century.^[21] In 1795, mahogany was introduced to India.

Phenology:

Soon after the new leaves develop, in April or May, the unassuming flowers appear. October through December is when the fruits ripen. It blooms in South India between February and May. Insects are the creatures that pollinate. Hybridization occurs often wherever the species coexist, particularly with *S. macrophylla*. Typically, only one flower in a cluster develops into a fruit after fertilization, while the others do not. It takes 8–10 months for a bloom to develop into a fully ripe fruit⁶. Fruits ripen between December and January. Since the fruit takes a long time to develop, crop evaluation can typically be done many months prior to harvest. Flowering typically occurs just before the rainy season, though it varies according to the climate, or geographic location. In the Caribbean Islands, the fruits of *S. mahagoni* Jacq. take eight to ten months to ripen, from January to March, after flowers bloom from April to July. Mahoganies typically flower and fruit every year between the ages of 10 and 15.^[23-26]



Figure 2: Fruit capsules of *Swietenia mahagoni* Jacq.

Growth & Reproduction: *Swietenia mahagoni* Jacq. is typically cultivated from seed because it is challenging to start from cuttings. The wind disperses the tiny, winged seeds of mahogany, which can produce a large number of seedlings close to mature trees. Direct sowing, transplanting, and stump planting are all methods of artificially propagating the tree; however, transplantation yields the best results and is the most widely used technique. Although pretreatment is usually not required, Immersing in water for 12 hours may improve the germination of seeds with low moisture content that have been kept. The seeds' ability to germinate is roughly 40%. The seeds are planted directly in containers or in holes or furrows that are 3–7 cm deep in a bed of light sand. One could say that seedlings fall somewhere in the middle of the light-demanding and shade-bearer spectrums in terms of their light requirements. Seeds that are germinating should be kept wet and under shade. It will take 10 to 21 days for seeds to sprout. Hypogeal germination occurs. Until they are out planted, the seedlings are maintained in shadow. When the seedlings are between 50 and 100 cm tall, they can be planted in the field.^[28]

Plant Collection Techniques: The capsules are harvested from the ground after the seeds fall or directly from trees between October and December, ideally just before they open. Site and year [5] have an impact on seed output. The seeds are removed and kept once the capsules open after being sun-dried. In India, seeds are stored for up to eight or even twelve months in a covered container. After six months, the feasibility of an open basket drops to a negligible level. The weight of the seeds varies. The seeds in Puerto Rico are said to weigh 7,040 seeds per kilogram, while the seeds in India weigh 3,350 to 3,500 seeds per kilogram. [30]



Figure 3: *S. mahogini jacq*

MATERIALS AND METHOD

Silver nitrate, swietenia mahagoni jacq Sodium potassium tartrate, starch, α -amylase enzyme, NaOH, NaCl₂, Sodium phosphate monophasic.

METHODOLOGY

Using the extraction methods like soxhlet extraction, benzene extraction, Ethanolic Extraction, Ethyl acetate extraction, petroleum ether extraction.⁶ Identification test of plant constituents parameters like flavonoids, terpenoids, saponins, carbohydrate and glycosidic compounds, proteins and amino acids, protein test, lignin. Biological assays, including in vitro and in vivo models, can be employed to identify novel compounds for the treatment of various diseases. Animals were randomly divided into 5 groups.

- Group 1- normal control,
- Group 2- diabetic control,
- Group3 –treatment control Swietenia mahagoni 100mg/kg,
- Group 4 – Swietenia mahagoni 200mg/kg treated group,
- Group 5 – Glibenclamide 5mg/kg.

The levels of biochemical parameter, body weight, serum lipid profile, serum total cholesterol, serum glycerides, VLDL AND HDL cholesterol, HDL cholesterol, Hemoglobin & glycated hemoglobin were examined in each of these groups. Chronic administration of swietenia amahagoni at a dose of 100mg/kg & 200mg/kg, p.o. for a period of 21 days markedly decreased the level of blood glucose level & lipid level. Pharmacological studies were assessed for bioassessment, clinical trials, and toxicological

studies.⁷And finally investigated quantitative estimation of total phenolic and flavonoid like estimation of total phenolic content, total flavonoid content, HPTLC investigation of ethanol extract. ⁸The following metrics were used to characterize Swieteniamahagoni Jacq. silver nanoparticles.

1. Visual examination,
2. UV- spectroscopy,
3. Fourier Transform Infrared Spectroscopy,
4. Drug Entrapment,
5. Determination of particle size & zeta potential, Scanning electron microscopy, &
6. In-vitro studys for first order kinetics, zero order kinetics, Higuchi's model, korsmayer peppas.

RESULTS AND DISCUSSION

A subset of the broader phrase "complementary and alternative medicine" (CAM) was another name for herbal medicine. Herbs were the standard treatment for almost all illnesses long before modern medicine was developed. Due to the negative consequences of contemporary medicine, more and more individuals are using herbal therapy as a supplement to and alternative to contemporary medications. The chemical components found in different plant tissues, referred to as secondary metabolites, and were what give medicinal plants their value. These compounds include alkaloids, glycosides, fatty and essential oils, gums, resins, mucilage, tannins, and more. The plant's storage organs, such as its roots, seeds, leaves, wood, etc., may contain these active ingredients.

Diabetes disrupts essential biochemical processes in nearly every cell and raises the chance of serious complications including atherosclerosis, neuropathy, cardiac infarction, and nephropathy, making it a major global health challenge. Therefore, one of the main areas of research has been the hunt for safer and more effective hypoglycemic medicines. The active plant components that were found to be a natural hypoglycemic medication due to traditional knowledge are used in many medical systems. Compared to synthetic medications, herbal ones were thought to have fewer negative effects.

They were more widely used, less harmful, and reasonably priced. The current study aims to determine which plant-based medication based on herbs may be utilized to treat diabetes mellitus, with a particular focus on assessing Swietenia Mahagoni Jacq. Ag nanoparticles that were mediated.

PHYTOCHEMICAL SCREENING

Preliminary screening tests on solvent extracts of *S. mahagoni Jacq*. Seeds revealed petroleum ether and benzene extracts contain alkaloids, carbohydrates, flavonoids, and proteins, while ethyl ethanoate extract lacks fixed oils and fats. All phytochemicals were found in the ethanolic fraction.

Determination of Phenolic and Flavonoid Levels

Using the standard curve, the phenolic content of *S. mahagoni* Jacq. seed extracts was quantified as µg gallic acid equivalents (GAE) per mg of sample. Table 9.3 & 9.4 provides absorbance value of gallic acid & Quercetin. The benchmark for flavonoid content was quercetin.




Figure 4: Calibration curve of Gallic acid.

Pharmacological studies: Pharmacological research was conducted in two distinct areas. The initial step was to use the α -amylase inhibitor test for in-vitro screening in arrange to conclude how effective the plant extracts were. The research's second component involved screening diabetic rats, OGGT, and in vivo.

In-vivo study

Acute oral toxicity studies: Oral Acute Toxicity of Ethanolic Extract of *S. mahagoni* Jacq. Seed was assessed according to OECD 425 guidelines. No mortality or toxic symptoms were observed within 14 days after administering a 2g/kg dose, indicating an LD₅₀ greater than 2 g/kg.

Oral glucose tolerance test: The study showed that *S. mahagoni* Jacq. seed extract enhanced glucose tolerance in rats, with 100 mg/kg being most effective. Consequently, the research evaluated its ethanol extract (0.1 and 0.2g/kg) for antioxidant, antidiabetic, and antihyperlipidemic effects in STZ-NC-induced diabetic rats.

Pre-formulation study

Solubility Test: The solubility test results for powdered *Swietenia mahagoni* Jacq seed extract in various solvents. Findings revealed that the powdered SMSE was readily soluble in ethanol and water, whereas it showed limited solubility in chloroform and diethyl ether. Consequently, the extract demonstrated greater solubility in polar solvents compared to non-polar ones.

UV-visible spectral analysis of *S.Mahagoni* Jacq. seed extract

A 0.05 mg sample of *S. mahagoni* Jacq. seed powder was dissolved in 10 mL purified water, and its UV-Vis spectrum (200–800 nm) is shown in Figure 9.26.




Figure 5: Uv-visible absorption spectra of S.Mahagoni Jacq. Seed extract

The greatest absorption of *S.Mahagoni* Jacq.wood and bark extract was discovered at 279nm, which was chosen as the wavelength for future investigation.

Calibration curve: UV-Visible Spectral Analysis of *S. mahagoni* Jacq Seed Extract at 279 nm. An R2 value of 0.999 indicated that the standard curve was linear between dosages ranging from 4 to 20 µg/ml. The sample *S. Mahagoni* Jacq. bark and wood aqueous extract, following Beer's law, has a concentration of 4-20µg/ml.

Green synthesis of Ag nano particles: Bark and wood extract from *S. Mahagoni* Jacq. were mixed with an Ag nitrate solution dropwise, and The Mixture was magnetically Processed at 120 rpm using hot plate to promote Ag nanoparticle synthesis. When Ag nanoparticles form, they turn brown instead of yellow.

Characterization of Ag nanoparticles contain *S. Mahagoni* ACQ. Extract

Visual Examination: When the Ag nitrate solution was added, the *S. Mahagoni* Jacq. extract's original yellow hue changed to a dark brown. The absence of a discernible color shift after ninety minutes indicated that the reduction process had finished. With a size in the nanometric range, This change suggested that the reaction medium's Ag ions had changed into elemental Ag.

UV Visible Spectral Analysis: The figure 2 showed periodic sampling that was done at 30-, 90-, 210-, and 24-hour intervals: 12. The study found that active biomolecules in the bark and wood extract of *Pterocarpus marsupium* Roxb. reduced Ag⁺ to Ag⁰, resulting in a color change from yellow to brown. The formation of silver nanoparticles, which

Table 1: Total Phenolic and Flavonoid Content of Extracts of *S. mahagoni* Jacq

S. No	Name of the Extract	Total Phenolic Content (mE GA/mg of dry extract)	Total Flavonoidal Content (mE quercetin/mg of dry extract)
1.	Pet ether extract	1.8 ± 0.14	3.55 ± 0.13
2.	Benzene extract	1.8 ± 0.07	1.3 ± 0.14
3.	Ethyl acetate extract	2.86 ± 0.10	2.2 ± 0.21
4.	Ethanol extract	2.96 ± 0.04	0.61 ± 0.03

Values are expressed as Mean ± SD (n = 3).

Table 2: Effect of ethanol extract of *S. mahagoni* seed on the body weight of rats at 2,000 mg/kg dose after 14 days

Group	Treatment	Body Weights (g) Before Treatment (M1 ± SD1)	Calculated 't' value After Treatment (M2 ± SD2)	Remarks
Control	Gum acacia suspension	160.63 ± 1.87	165.06 ± 2.08	t = 2.7412
Treated	2,000 mg/kg of EESM (Ethanol Extract of <i>S. mahagoni</i>)	163.03 ± 3.40	167.50 ± 2.52	t = 1.8264

Table 3: Effect of Ethanol extract of *S. Mahagoni* Jacq seed on oral glucose tolerance test (OGTT) in normal healthy rats

Group	Treatment	Dose (mg/kg)	Fasting blood glucose level (mg/dL) 0 min	Reduction after 60 min (%) 30 min
I	Normal Control	----	72.33 ± 1.23	106.00 ± 1.94
II	Animal 1	100	82.00 ± 1.80	112.00 ± 1.90
III	Animal 2	200	81.54 ± 1.40	110.00 ± 1.16
IV	Animal 3	300	78.00 ± 1.40	107.32 ± 1.60
VIII	Glibenclamide	5	79.50 ± 1.87	106.33 ± 1.70

Table 4: Effect of ethanolic seed extract of *S. mahagoni* Jacq. on HDL, LDL, TC, TG of control and experimental groups of rats

GROUP	HDL	LDL	TC	TG
Normal control	52.83 ± 1.014	23.33 ± 0.88	129.01 ± 1.065	63.51 ± 1.66
Diabetic control	13.33 ± 1.909###	145.7 ± 2.04###	280.2 ± 0.75###	201.88 ± 3.911###
STZ + G	42.17 ± 0.703**	39.67 ± 1.22**	156.54 ± 2.232**	90.52 ± 2.203**
STZ SM 100 mg	26.00 ± 1.06*	74.83 ± 0.70**	217.52 ± 0.45**	182.00 ± 2.95**
STZ SM 200 mg	37.17 ± 0.70**	53.17 ± 0.60**	177.53 ± 0.30**	134.83 ± 3.148**

The data are expressed as mean ± SEM, n = 6.

One-way ANOVA followed by Dunnett's test. ns = non-significant
Compared with normal control: p < 0.05, ## p < 0.01, ### p < 0.001
Compared with disease control: p < 0.05, ** p < 0.01, *** p < 0.001

Table 5: Solubility of *S. Mahagoni* Jacq. in different solvents

Solvent	Soluble	Sparingly soluble	Insoluble
Ethanol	+	-	-
Water	+	-	-
Chloroform	-	+	-
Diethyl ether	-	-	-

display a surface plasmon resonance (SPR) absorption band in the visible spectrum, is linked to this transformation.

Electrostatic interactions including hydrogen bonds, bio-organic connections, and capping molecules are what lead to the biogenesis of Ag nanoparticles. Additional information on morphology can be obtained by scanning electron microscopy. It was discovered from the SEM examination of *S. Mahagoni* Jacq. Ag nanoparticles are spherical in form and assemble on the surface.¹⁰

Histopathological studies: At the conclusion of the investigation, the pancreatic histology of the experimental rats was assessed.^{9,10} Diabetes-induced pancreas exhibited severe necrotic alterations, reduced β cell size, and diminished Langerhans Islets. However, rats treated with STZ-NC showed normal β cell size, acini, and Islets of Langerhans, restoring normal structures.




Figure 6: Uv-visible absorption spectra of silver nanoparticles at different intervals




Figure 7: SEM analysis of S. Mahagoni Jacq Seed extract nanoparticles at 1000X magnification




Figure 8: Effect of ethanolic seed extract of S. mahagoni Jacq. on cellular damage in pancreas of control and experimental groups of rats (400x)

B = Diabetic control group (showing atrophic acini, reduced β -cell size, and decreased islet size); C = STZ + Glibenclamide group (demonstrating well-regenerated and preserved cells with significant β -cell proliferation and regeneration); E=STZ+SM 200mg group (exhibiting Functional, regenerated, and conserved cells with notable β -cell proliferation and regeneration); A and D=STZ+SM 100mg group (displaying Pancreas exhibiting atrophy with acinar atrophy, reduced β -cell size, and presence of hyperplastic cells).

Statistical Analysis: The study used Graph Pad Prism 5 for statistical analysis of differences and

linear regression, expressing all values as mean $\pm$ SD.

CONCLUSION

Since the beginning of the world civilization humans are depended on herbal medicine because of its health care and various disease treatment properties. Over 300 limonoids have been extracted from this family, including seven from the extract of *S. mahagoni* seeds. Different species of *Swietenia* contain in their various parts phenolic compounds, triterpenoids (limonoids), flavonoids, swiemahogins A and BC, and alkaloids. The chemical entities of this plant have been proved for their Anti-bacterial activity, Antimicrobial Activity, Anti-oxidant activity, Antiulcer activity, Anti-fungal activity, Anti inflammatory, Analgesic activity, Hypoglycemic activity, Platelet Aggregation Inhibitors activity etc. These scientifically proved activities can be related with the traditional usage of the plant. Thus *S. mahagoni* Jacq. is one of the most important plants that has a tremendous scope for research in future. The novelty and applicability of this valuable species are hidden. Such things should be overcome through extensive scientific research. The drug may be a good candidate for developing a safe, tolerable, and promising nutraceutical treatment for the management of many diseases.

REFERENCES

1. Adewole, S. O., & Ojewole, J. A. O. (2007). Insulin-induced immunohistochemical and morphological changes in pancreatic β -cells of streptozotocin-treated diabetic rats. *Methods and Findings in Experimental and Clinical Pharmacology*, 29(7), 447–455.
2. Ahmad, A. R., Malik, A., & Handayani, V. (2023). *In vivo* and *in vitro* antidiabetic assay of purified mahoni seeds extract (*Swietenia mahagoni* (L.) Jacq.). *Biomedical and Pharmacology Journal*, 16(3), 1701–1706. <https://doi.org/10.13005/bpj/2748>
3. Ahamd, H., et al. (2020). The recent use of *Swietenia mahagoni* (L.) Jacq. as antidiabetes type 2 phytomedicine: A systematic review. *Heliyon*, 6(3), e03536. <https://doi.org/10.1016/j.heliyon.2020.e03536>
4. Buzea, C., Pacheco, I. I., & Robbie, K. (2007). Nanomaterials and nanoparticles: Sources and toxicity. *Biointerphases*, 2(4), MR17–MR71. <https://doi.org/10.1116/1.2815690>
5. Cruze, L. R. M. D., Shanta, A. A., Mostaq, M. S., Mostafa, A., Hasan, M. K., Rashid, F., & Afrin, S. (2020). An *in vivo* evaluation of antidiabetic and depressant action of bark of *Swietenia mahagoni* on alloxan-induced diabetic rat model. *Asian Journal of Research and Reports in Endocrinology*, 3(1), 71–79. <https://doi.org/10.9734/ajrre/2020/v3i130>
6. Darussalam, H., Nuryastuti, T., Mursiti, S., & Mustofa, M. (2014). Antimicrobial activity of bioactive compounds isolated from *Swietenia mahagoni*

- (L.) Jacq. against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *Journal of the Medical Sciences*, 46(4), 182–188. <https://doi.org/10.19106/JMedScie004604201402>
7. Dhir, R., Chauhan, S., Subham, P., Kumar, S., Sharma, P., Shidiki, A., & Kumar, G. (2024). Plant-mediated synthesis of silver nanoparticles: Unlocking their pharmacological potential—A comprehensive review. *Frontiers in Bioengineering and Biotechnology*, 11, 1324805. <https://doi.org/10.3389/fbioe.2023.1324805>
8. Ghosh, S., Besra, S. E., Roy, K., Gupta, J. K., & Vadasiramon, J. R. (2010). Pharmacological effects of methanolic extract of *Swietenia mahagoni* Jacq. (Meliaceae) seeds. *International Journal of Green Pharmacy*, 4(2), 115–120.
9. Hammam, M. A., Kalil, G. A., El-Sayed, S. M., & Mohamed, I. I. (2020). *Swietenia mahagoni* seeds attenuate hyperglycemia and protect liver in alloxan-induced diabetic rats. *Menoufia Journal of Agricultural Biotechnology*, 5(4), 103–112. <https://doi.org/10.21608/mjab.2020.171468>
10. Heidegger, S., Niedermayer, S., Schmidt, A., Gößl, D., Argyo, C., Endres, S., Bein, T., & Bourquin, C. (2015). Immune response to functionalized mesoporous silica nanoparticles for targeted drug delivery. *arXiv preprint*. <https://arxiv.org/abs/1509.01029>
11. Liu, J., Peng, X., Zhang, W., Shi, L., Li, X., Chen, J., & Qiu, M. (2013). Swietemahalactone, a rearranged phragmalin-type limonoid with antibacterial effect, from *Swietenia mahagoni*. *RSC Advances*, 3(12), 4890–4893. <https://doi.org/10.1039/C3RA23401K>
12. Mahid-Al-Hasan, S. M. M., Khan, M. I., & Umar, B. U. (2011). Effect of ethanolic extract of *Swietenia mahagoni* seeds on experimentally induced diabetes mellitus in rats. *Faridpur Medical College Journal*, 6(2), 70–73. <https://doi.org/10.3329/fmcj.v6i2.9203>
13. Maurya, A., & Tyagi, S. (2024). Recent advances in nanoparticle-based drug delivery systems. *Journal of Drug Discovery and Health Sciences*, 1(4), 201–211. <https://doi.org/10.21590/jddhs.01.04.03>
14. Muthu, K., Akilandaewaswari, B., & Nagasundari, S. M. (2019). Green synthesis of silver nanoparticles using *Polyalthia longifolia* stem bark extract and its catalytic reduction of 4-nitrophenol. *Asian Journal of Chemistry*, 31(11), 2439–2442. <https://doi.org/10.14233/ajchem.2019.22071>
15. Nkosi, C., Basson, A. K., Ntombela, Z. G., Dlamini, N. G., & Pullabhotla, R. V. S. R. (2024). Green synthesis, characterization, and application of silver nanoparticles using bioflocculant: A review. *Bioengineering*, 11(5), 492. <https://doi.org/10.3390/bioengineering11050492>
16. Panda, S. S., Jafri, M., Kar, A., & Meheta, B. K. (2010). Antidiabetic potentiality of the aqueous-methanolic extract of seed of *Swietenia mahagoni* (L.) Jacq. in streptozotocin-induced diabetic male albino rat. *Journal of Ethnopharmacology*, 129(2), 239–246. <https://doi.org/10.1016/j.jep.2010.03.013>
17. Pratiwi, S., Emelda, K., Munir, M. A., Azizah, A., & Pothitirat, W. (2024). Analysis of total phenolic content and antioxidant activity of mahogany seed infusion (*Swietenia mahagoni* (L.) Jacq.). *Indonesian Journal of Nutrition and Dietetics*, 12(4), 295–304.
18. Ranjith, S., & Jeevanandam, M. (2024). A review on green synthesized metal nanoparticles applications. *Journal for Research in Applied Sciences and Biotechnology*, 3(5), 80–100. <https://doi.org/10.55544/jrasb.3.5.11>
19. Shahzadi, S., Fatima, S., Ul Ain, Q., Shafiq, Z., & Janjua, M. R. S. A. (2025). A review on green synthesis of silver nanoparticles (SNPs) using plant extracts: A multifaceted approach in photocatalysis, environmental remediation, and biomedicine. *RSC Advances*, 15, 3858–3903. <https://doi.org/10.1039/D4RA07519F>
20. Shidiki, A., & Kumar, G. (2023). Green synthesis of silver nanoparticles: Methods, biological applications, delivery, and toxicity. *Materials Advances*, 4(4), 1831–1849. <https://doi.org/10.1039/D2MA01105K>
21. Stephan, R., Jeyabharathi, S., Jeenathunisa, N., & Deena Priscilla, H. (2018). Antidiabetic activity of Indian mahagoni plant extract by transdermal therapy. *Asian Journal of Pharmaceutical and Clinical Research*, 11(4), 361–365. <https://doi.org/10.22159/ajpcr.2018.v11i4.23234>
22. Sukardiman, S., et al. (2018). Normalization of insulin resistance, glucose intolerance, and lipid profile by *Swietenia mahagoni* leaf extract in fructose-induced diabetic rats. *Pharmacognosy Magazine*, 14(59), 349–357. <https://doi.org/10.4103/pm.pm.358.18>
23. Villaseñor-Cavazos, F. J., Torres-Valladares, D., & Lozano, O. (2021). Modelling and optimization of nanovector synthesis for applications in drug delivery systems. *arXiv preprint*. <https://arxiv.org/abs/2112.02002>
24. Vidyasagar, R., Patel, R. R., Singh, S. K., & Singh, M. (2023). Green synthesis of silver nanoparticles: Methods, biological applications, delivery, and toxicity. *Materials Advances*, 4(6), 1831–1849.
25. Widayati, T., & Pertiwi, A. D. (2024). Phytochemical profile and potential of *Swietenia mahagoni* as a source of bioactive compounds. *Pharmacognosy Reviews*, 18(36), 112–121.
26. Wulandari, D. E., & Arifin, Z. (2024). Preclinical safety evaluation of *Swietenia mahagoni* leaf in

- Wistar rats. *International Journal of Pharmacy and Pharmaceutical Sciences*, 16(3), 89–94.
27. Zakaria, Z., & Nabila, A. (2023). Normalization of lipid profile by *Swietenia mahagoni* seed extract in diabetic rats. *Pharmacology & Toxicology Reports*, 12(2), 41–47.
 28. Zhang, Y., Li, C., & Wang, X. (2024). Nanomedicine for targeted drug delivery systems: A mini review. *International Journal of Nanotechnology and Allied Sciences*, 8(2), 29–36.
 29. Zhao, L., Chen, Y., & Wang, Z. (2025). Recent updates on green synthesized silver nanoparticles: Preparation technologies, properties, and biomedical applications. *International Journal of Environmental Sciences*, 11(13s), 968–982.
 30. Zohra, F., Khan, S., & Ahmad, N. (2018). Green synthesis of silver nanoparticles from *Tectona grandis* seed extract: Characterization and mechanism of antimicrobial action. *Journal of Analytical Science and Technology*, 9, 13.
<https://doi.org/10.1186/s40543-018-0163-z>